

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Draxxin 25 mg/ml solution for injection for pigs

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

#### **Active substance:**

Tulathromycin 25 mg/ml

#### **Excipient:**

Monothioglycerol 5 mg/ml

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Solution for injection.

Clear colourless to slightly yellow solution.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Pigs.

#### **4.2 Indications for use, specifying the target species**

Treatment and metaphylaxis of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Mycoplasma hyopneumoniae*, *Haemophilus parasuis* and *Bordetella bronchiseptica* susceptible to tulathromycin. The presence of the disease in the group must be established before the product is used. The veterinary medicinal product should only be used if pigs are expected to develop the disease within 2–3 days.

#### **4.3 Contraindications**

Do not use in cases of hypersensitivity to macrolide antibiotics or to any of the excipients.

#### 4.4 Special warnings for each target species

Cross resistance occurs with other macrolides. Do not administer simultaneously with antimicrobials with a similar mode of action such as other macrolides or lincosamides.

#### 4.5 Special precautions for use

##### Special precautions for use in animals

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria. Official, national and regional antimicrobial policies should be taken into account when the product is used. Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to tulathromycin and may decrease the effectiveness of treatment with other macrolides, lincosamides and group B streptogramins, due to the potential for cross resistance.

If a hypersensitivity reaction occurs appropriate treatment should be administered without delay.

##### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Tulathromycin is irritating to eyes. In case of accidental eye exposure, flush the eyes immediately with clean water.

Tulathromycin may cause sensitisation by skin contact resulting in e.g. reddening of the skin (erythema) and/or dermatitis. In case of accidental spillage onto skin, wash the skin immediately with soap and water.

Wash hands after use.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

If there is suspicion of a hypersensitivity reaction following accidental exposure (recognised by e.g. itching, difficulty in breathing, hives, swelling on the face, nausea, vomitus) appropriate treatment should be administered. Seek medical advice immediately and show the package leaflet or the label to the physician.

#### 4.6 Adverse reactions (frequency and seriousness)

Pathomorphological injection site reactions (including reversible changes of congestion, oedema, fibrosis and haemorrhage) are present for approximately 30 days after injection.

#### 4.7 Use during pregnancy, lactation or lay

Laboratory studies in rats and rabbits have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. The safety of the veterinary medicinal

product has not been established during pregnancy and lactation. Use only according to the benefit/risk assessment by the responsible veterinarian.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

None known.

#### **4.9 Amounts to be administered and administration route**

A single intramuscular injection of 2.5 mg tulathromycin/kg bodyweight (equivalent to 1 ml/10 kg bodyweight) in the neck.

For treatment of pigs over 40 kg bodyweight, divide the dose so that no more than 4 ml are injected at one site.

For any respiratory disease, it is recommended to treat animals in the early stages of the disease and to evaluate the response to treatment within 48 hours after injection. If clinical signs of respiratory disease persist or increase, or if relapse occurs, treatment should be changed, using another antibiotic, and continued until clinical signs have resolved.

To ensure correct dosage bodyweight should be determined as accurately as possible to avoid underdosing. For multiple vial entry, an aspirating needle or multi-dose syringe is recommended to avoid excessive broaching of the stopper.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In young pigs weighing approximately 10 kg given three or five times the therapeutic dose transient signs attributed to injection site discomfort were observed and included excessive vocalisation and restlessness. Lameness was also observed when the hind leg was used as the injection site.

#### **4.11 Withdrawal period(s)**

Meat and offal: 13 days.

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: Antibacterials for systemic use, macrolides.  
ATCvet code: QJ01FA94.

#### **5.1 Pharmacodynamic properties**

Tulathromycin is a semi-synthetic macrolide antimicrobial agent, which originates from a fermentation product. It differs from many other macrolides in that it has a long duration of action that is, in part, due to its three amine groups; therefore it has been given the chemical subclass designation of triamilide.

Macrolides are bacteriostatic acting antibiotics and inhibit essential protein biosynthesis by virtue of their selective binding to bacterial ribosomal RNA. They act

by stimulating the dissociation of peptidyl-tRNA from the ribosome during the translocation process.

Tulathromycin possesses *in vitro* activity against *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Mycoplasma hyopneumoniae*, *Haemophilus parasuis* and *Bordetella bronchiseptica*, the bacterial pathogens most commonly associated with swine respiratory disease. Increased minimum inhibitory concentration (MIC) values have been found in some isolates of *Actinobacillus pleuropneumoniae*.

The Clinical and Laboratory Standards Institute CLSI has set the clinical breakpoints for tulathromycin against *P. multocida* and *B. bronchiseptica* of swine respiratory origin, as  $\leq 16$   $\mu\text{g/ml}$  susceptible and  $\geq 64$   $\mu\text{g/ml}$  resistant. For *A. pleuropneumoniae* of swine respiratory origin the susceptible breakpoint is set at  $\leq 64$   $\mu\text{g/ml}$ . CLSI has also published clinical breakpoints for tulathromycin based on a disk diffusion method (CLSI document VET08, 4th ed, 2018). No clinical breakpoints have been set for *H. parasuis*. Neither EUCAST nor CLSI have developed standard methods for testing antibacterial agents against veterinary *Mycoplasma* species and thus no interpretative criteria have been set.

Resistance to macrolides can develop by mutations in genes encoding ribosomal RNA (rRNA) or some ribosomal proteins; by enzymatic modification (methylation) of the 23S rRNA target site, generally giving rise to cross-resistance with lincosamides and group B streptogramins (MLS<sub>B</sub> resistance); by enzymatic inactivation; or by macrolide efflux. MLS<sub>B</sub> resistance may be constitutive or inducible. Resistance may be chromosomal or plasmid-encoded and may be transferable if associated with transposons, plasmids, integrative and conjugative elements. Additionally, the genomic plasticity of *Mycoplasma* is enhanced by the horizontal transfer of large chromosomal fragments.

In addition to its antimicrobial properties, tulathromycin demonstrates immunomodulating and anti-inflammatory actions in experimental studies. In porcine polymorphonuclear cells (PMNs; neutrophils), tulathromycin promotes apoptosis (programmed cell death) and the clearance of apoptotic cells by macrophages. It lowers the production of the pro-inflammatory mediators leukotriene B<sub>4</sub> and CXCL-8 and induces the production of anti-inflammatory and pro-resolving lipid lipoxin A<sub>4</sub>.

## 5.2 Pharmacokinetic particulars

In pigs, the pharmacokinetic profile of tulathromycin when administered as a single intramuscular dose of 2.5 mg/kg bodyweight, was also characterised by rapid and extensive absorption followed by high distribution and slow elimination. The maximum concentration ( $C_{\text{max}}$ ) in plasma was approximately 0.6  $\mu\text{g/ml}$ ; this was achieved approximately 30 minutes post-dosing ( $T_{\text{max}}$ ). Tulathromycin concentrations in lung homogenate were considerably higher than those in plasma. There is strong evidence of substantial accumulation of tulathromycin in neutrophils and alveolar macrophages. However, the *in vivo* concentration of tulathromycin at the infection site of the lung is not known. Peak concentrations were followed by a slow decline in systemic exposure with an apparent elimination half-life ( $t_{1/2}$ ) of approximately 91 hours in plasma. Plasma protein binding was low, approximately 40%. The volume of distribution at steady-state ( $V_{\text{ss}}$ ) determined after intravenous administration was 13.2 L/kg. The bioavailability of tulathromycin after intramuscular administration in pigs was approximately 88%.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Monothioglycerol  
Propylene glycol  
Citric acid  
Hydrochloric acid  
Sodium hydroxide  
Water for injections

### **6.2 Major incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.  
Shelf life after first opening the immediate packaging: 28 days.

### **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

### **6.5 Nature and composition of immediate packaging**

Primary packaging: Type I glass vial with a fluoropolymer coated chlorobutyl stopper and an aluminium overseal.

Pack size: Cardboard box containing one vial.

Vial sizes: 50, 100 and 250 ml.

Not all pack sizes may be marketed.

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP

**8. MARKETING AUTHORISATION NUMBER**

Vm 42058/5022

**9. DATE OF FIRST AUTHORISATION**

8 July 2014

**10. DATE OF REVISION OF THE TEXT**

August 2021

Approved 24 August 2021

