

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Marbocare flavour 5mg tablets for dogs and cats

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

**Each tablet contains:**

**Active substance:** Marbofloxacin 5.0 mg

**Excipients:**

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Tablet

Beige brown spotted round tablets with a score line on one side.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Dogs and Cats

#### **4.2 Indications for use, specifying the target species**

Marbofloxacin is indicated in the treatment of the following infections caused by susceptible strains of organisms (see section 5.1);

##### Dogs

- Skin and soft tissue infections (skinfold pyoderma, impetigo, folliculitis, furunculosis, cellulitis)
- Urinary tract infections (UTI) associated or not with prostatitis or epididymitis
- Respiratory tract infections

##### Cats

- Skin and soft tissue infections (wounds, abscesses, phlegmons)
- Upper respiratory tract infections

#### **4.3 Contraindications**

Do not use in dogs aged less than 12 months, or less than 18 months for exceptionally large breeds of dogs, such as Great Danes, Briard, Bernese, Bouvier and Mastiffs, with a longer growth period.

Do not use in cats aged less than 16 weeks.

Do not use in animals with known hypersensitivity to marbofloxacin or other (fluoro)quinolones or to any of the excipients.

Do not use in cases of confirmed or suspected resistance to fluoroquinolones (cross resistance).

#### **4.4 Special warnings for each target species**

A low urinary pH could have an inhibitory effect on the activity of marbofloxacin.

#### **4.5 Special precautions for use**

##### **Special precautions for use in animals**

The fluoroquinolones have been shown to induce erosion of articular cartilage in juvenile dogs and care should be taken to dose accurately especially in young animals. However, at the therapeutic recommended dosage, no severe side effects are to be expected in dogs and cats.

Some fluoroquinolones at high doses may have an epileptogenic potential. Cautious use is recommended in dogs and cats diagnosed as suffering from epilepsy.

Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly to other classes of antimicrobials. Whenever possible, use of fluoroquinolones should be based on susceptibility testing. Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to the fluoroquinolones and may decrease effectiveness of treatment with other quinolones due to the potential for cross-resistance. Official and local antimicrobial policies should be taken into account when the product is used.

##### **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

People with known hypersensitivity to (fluoro)quinolones should avoid using this product. Avoid contact of the skin and eyes with the product. In case of accidental ingestion seek medical advice immediately and show the package leaflet or the label to the physician. Wash hands after use.

#### **4.6 Adverse reactions (frequency and seriousness)**

Mild side effects such as vomiting, softening of faeces, modification of thirst or transient increase in activity may very rarely occur. These signs cease spontaneously after treatment and do not necessitate cessation of treatment.

The frequency of adverse reactions is defined using the following convention:

- Very common (more than 1 in 10 animals treated displaying adverse reaction(s))

- Common (more than 1 but less than 10 animals in 100 animals treated)
- Uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- Rare (more than 1 but less than 10 animals in 10,000 animals treated)
- Very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

#### 4.7 Use during pregnancy, lactation or lay

Studies in laboratory animals (rats, rabbits) showed no teratogenicity, embryotoxicity and maternotoxicity with marbofloxacin at therapeutic doses.

The safety of marbofloxacin has not been assessed in pregnant and lactating cats and dogs. Use only accordingly to the benefit/risk assessment by the responsible veterinarian in pregnant and lactating animals.

#### 4.8 Interaction with other medicinal products and other forms of interaction

Fluoroquinolones are known to interact with orally administered cations (Aluminium, Calcium, Magnesium, Iron). In such cases, the bioavailability may be reduced.

When administered together with theophylline, the half-life and thus the plasma concentration of theophylline increase. Hence, in case of concurrent administration the dose of theophylline should be reduced.

Do not use in combination with tetracyclines, macrolides because of the potential antagonist effect.

#### 4.9 Amounts to be administered and administration route

For oral administration.

The recommended dose rate is 2 mg/kg/day in a single daily administration (see table below).

Body Weight	Tablets
1.3 – 2.5 kg	1
2.6 – 5.0 kg	2

To ensure a correct dosage body weight should be determined as accurately as possible to avoid underdosing.

##### DOGS:

- In skin and soft tissue infections, treatment duration is at least 5 days. Depending on the course of the disease, it may be extended up to 40 days.
- In urinary tract infections, treatment duration is at least 10 days. Depending on the course of the disease, it may be extended up to 28 days.
- In respiratory infections, treatment duration is at least 7 days and depending on the course of the disease, it may be extended up to 21 days.

CATS:

- For skin and soft tissue infections (wounds, abscesses, phlegmons) treatment duration is 3 to 5 days.
- For upper respiratory infections treatment duration is 5 days.

#### 4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Overdosage may cause acute signs in the form of neurological disorders, which should be treated symptomatically.

#### 4.11 Withdrawal period(s)

Not applicable.

### 5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Antibacterials for systemic use;  
Fluoroquinolones; Marbofloxacin.  
ATC vet code: QJ01MA93

#### 5.1 Pharmacodynamic properties

Marbofloxacin is a synthetic, bactericidal antimicrobial, belonging to the fluoroquinolone group which acts by inhibition of DNA gyrase. It is effective against a wide range of Gram positive bacteria (in particular *Staphylococci*, *Streptococci*) and Gram negative bacteria (*Escherichia coli*, *Salmonella typhimurium*, *Citrobacter freundii*, *Enterobacter cloacae*, *Serratia marcescens*, *Morganella morganii*, *Proteus spp*, *Klebsiella spp*, *Shigella spp*, *Pasteurella spp*, *Haemophilus spp*, *Moraxella spp*, *Pseudomonas spp*, *Brucella canis*) as well as *Mycoplasma spp*.

Bacterial strains with a MIC  $\leq$  1  $\mu$ g/ml are susceptible, strains with a MIC of 2  $\mu$ g/ml are intermediately susceptible and strains with a MIC  $\geq$  4  $\mu$ g/ml are resistant to marbofloxacin (CLSI, 2004).

Resistance to fluoroquinolones occurs mostly by chromosomal mutation with three mechanisms: decrease of the bacterial wall permeability, expression of efflux pump or mutation of enzymes responsible for molecule binding.

Marbofloxacin is not active against anaerobes, yeasts or fungi.

#### 5.2 Pharmacokinetic particulars

After oral administration in dogs and cats at the recommended dose of 2 mg/kg, marbofloxacin is readily absorbed and reaches maximal plasma concentrations of 1.5  $\mu$ g/ml within 2 hours

Its bioavailability is close to 100%.

It is weakly bound to plasma proteins (less than 10%), extensively distributed and in most tissues (liver, kidney, skin, lung, bladder, digestive tract) it achieves higher

concentrations than in plasma. Marbofloxacin is eliminated slowly ( $t_{1/2\beta} = 14$  h in dogs and 10h in cats) predominantly in the active form in urine (2/3) and faeces (1/3).

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose monohydrate  
Povidone (K90)  
Silica, colloidal hydrated  
Crospovidone (Type A)  
Castor Oil, Hydrogenated  
Dessicated pork liver powder  
Dried yeast  
Magnesium stearate

### **6.2 Major incompatibilities**

Not applicable.

### **6.3 Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.

### **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

### **6.5 Nature and composition of immediate packaging**

The product is packaged in Aluminium- PVC/aluminium/polyamide blister  
Box containing 1 blister of 10 tablets (10 tablets)  
Box containing 2 blisters of 10 tablets (20 tablets)  
Box containing 10 blisters of 10 tablets (100 tablets)

Not all pack sizes may be marketed.

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Emdoka Bvba

John Lijssentstraat 16  
B-2321 Hoogstraten  
Belgium

**8. MARKETING AUTHORISATION NUMBER**


Vm 34534/4003

**9. DATE OF FIRST AUTHORISATION**

04 September 2013

**10. DATE OF REVISION OF THE TEXT**

June 2020



Approved: 25 June 2020