

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Metacam 15 mg/ml oral suspension for pigs

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One ml contains:

Active substance(s):

Meloxicam 15 mg

Excipient(s):

Sodium benzoate 1.5 mg

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Oral suspension

Yellowish viscous oral suspension with a green tinge.

4. CLINICAL PARTICULARS

4.1 Target species

Pigs

4.2 Indications for use, specifying the target species

For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation. For adjunctive therapy in the treatment of puerperal septicaemia and toxemia (Mastitis-Metritis- Agalactia syndrome MMA) with appropriate antibiotic therapy.

4.3 Contraindications

Do not use in pigs suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

4.4 Special warnings for each target species

None.

4.5 Special precautions for use

Special precautions for use in animals

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

Avoid use in very severely dehydrated, hypovolaemic or hypotensive pigs which require parenteral rehydration, as there may be a potential risk of renal toxicity.

Special precautions to be taken by the person administering the veterinary medicinal product to animals People with known hypersensitivity to non-steroidal anti-inflammatory drugs (NSAIDs) should avoid contact with the veterinary medicinal product.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

This product can cause eye irritation. In case of contact with the eyes, immediately rinse thoroughly with water.

4.6 Adverse reactions (frequency and seriousness)

None.

4.7 Use during pregnancy, lactation or lay

Can be used during pregnancy and lactation.

4.8 Interaction with other medicinal products and other forms of interaction

Do not administer concurrently with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anticoagulant agents.

4.9 Amounts to be administered and administration route

Oral suspension to be administered at a dosage of 0.4 mg/kg body weight (i.e. 2.7 ml/100 kg) in combination with antibiotic therapy, as appropriate. If required, a second administration of Meloxicam can be given after 24 hours. In cases of MMA with severely disturbed general demeanour (e.g. anorexia) the use of Metacam 20 mg/ml solution for injection is recommended.

To be administered preferably mixed with a small quantity of feed. Alternatively to be given prior to feeding, or directly into the mouth.

The suspension should be given using the measuring syringe provided in the package. The syringe fits onto the bottle and has a kg-body weight scale.

Shake well before use.

After administration of the veterinary medicinal product, close the bottle by replacing the cap, wash the measuring syringe with warm water and let it dry.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

In case of overdose symptomatic treatment should be initiated.

4.11 Withdrawal period(s)

Meat and offal: 5 days.

5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Anti-inflammatory and antirheumatic products, non-steroids (oxicams). ATCvet code: QM01AC06.

5.1 Pharmacodynamic properties

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, analgesic, anti-exudative and antipyretic effects. It reduces leukocyte infiltration into the inflamed tissue. To a minor extent it also inhibits collagen-induced thrombocyte aggregation. Meloxicam also has anti-endotoxic properties because it has been shown to inhibit production of thromboxane B₂ induced by intravenous *E. coli* endotoxin administration in pigs.

5.2 Pharmacokinetic particulars

Absorption

After a single oral dose of 0.4 mg meloxicam/kg a C_{max} value of 0.81 µg/ml was reached after 2 hours.

Distribution

More than 98 % of meloxicam is bound to plasma proteins. The highest meloxicam concentrations are to be found in liver and kidney. Comparatively low concentrations are detectable in skeletal muscle and fat.

Metabolism

Meloxicam is predominantly found in plasma. Bile and urine contain only traces of the parent compound. Meloxicam is metabolised to an alcohol, an acid derivative and to several polar metabolites. All major metabolites have been shown to be pharmacologically inactive.

Elimination

After oral administration the mean plasma elimination half-life is approximately 2.3 hours. Approximately 50 % of the administered dose is eliminated via urine and the remainder via faeces.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium benzoate Sorbitol, liquid Glycerol Saccharin sodium
Xylitol
Sodium dihydrogen phosphate dihydrate Silica, colloidal
anhydrous Hydroxyethylcellulose
Citric acid Honey aroma Water, purified

6.2 Major incompatibilities

None known.

6.3 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 3 years. Shelf life after first opening of the immediate packaging: 6 months.

6.4 Special precautions for storage

This veterinary medicinal product does not require any special storage conditions.

6.5 Nature and composition of immediate packaging

Cardboard box containing one polyethylene bottle of 100 ml or 250 ml with a polyethylene tip adapter, a tamper-proof child-resistant closure and a measuring syringe. Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Boehringer Ingelheim
Vetmedica GmbH 55216
Ingelheim/Rhein
GERMANY

8. MARKETING AUTHORISATION NUMBERS

EU/2/97/004/041 100 ml
EU/2/97/004/042 250 ml

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 07.01.1998
Date of last renewal: 06.12.2007

10. DATE OF REVISION OF THE TEXT

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>

PROHIBITION OF SALE, SUPPLY AND/OR USE

Not applicable.