



Children's Chesty Coughs

Guaifenesin

6 Years +

- This medicine is used to help relieve children's chesty coughs.
- This medicine is for use by children aged 6-12 years.
- **Do not use this medicine:**
 - There are some people who should not use this medicine. *To find out if your child is one of them see section 2 ►*
 - If your child has ever had a **bad reaction** to any of the ingredients. *For the list of ingredients see section 6 ►*
- **Speak to your doctor:**
 - If your child suffers from any of the conditions mentioned in section 2. *See section 2 ►*
 - If you are taking any other medicines. *See section 2 ►*
- **Follow the dosage instructions carefully.** Some children should not take this product. Follow the instructions shown in the table carefully. *See section 3 ►*

Now read this whole leaflet carefully before you use this medicine. Keep the leaflet: you might need it again.

1 What the medicine is and what it is used for

Benylin Children's Chesty Coughs is a medicine which is used to help relieve chesty coughs in children. It contains guaifenesin (an expectorant) to help loosen mucus (phlegm) from the lungs and make it easier to cough up. The medicine is for use in children aged 6-12 years.

2 What you need to know before giving this medicine to your child

This medicine is suitable for most children but a few children should not use it. If you are in any doubt, talk to your doctor or pharmacist.

X Do not use this medicine...

- If your child has ever had a **bad reaction** to this product or any of the ingredients. *See section 6 for full list of ingredients.*
- If your child is taking **any other cough and cold medicine.**
- If your child is aged under 6 years.

If any of these apply, **get advice from a doctor or pharmacist without using Benylin Children's Chesty Coughs.**

! Talk to your doctor or pharmacist...

- If your child suffers from **liver or kidney problems.**
- If your child has a **persistent cough** such as occurs with asthma, has **asthma**, is suffering from an **asthma attack** or a **cough that produces a lot of mucus (phlegm).**

If any of these apply now or in the past, **talk to a doctor or pharmacist.**

! If you are pregnant or breast-feeding

- The following advice is included in case an older child or adult is taking the medicine.
- If you are pregnant or breast-feeding, only use this medicine on the advice of your doctor.

! Some of the ingredients can cause problems

- This medicine contains 5.05 g sorbitol in each 10 ml. Sorbitol is a source of fructose. If your doctor has told you that you (or your child) have an intolerance to some sugars or if you have been

diagn
(HFI)
canno
befor
medi
disco
■ This
per l
reco
sodiu
■ This
each
■ This
each
react
advic
you h
large
in yo
"met

3 How to use this medicine

- For oral use
- Do not give

i Child

This medicine is for use in children aged 6 years old.



i Children 6 - 12 years

Age	Dose
Children 6 – 12 years	Two 5 ml spoonfuls four times a day.

- Do not give more than 4 doses in 24 hours.
- If symptoms persist or worsen, talk to your doctor or pharmacist.
- Do not give to a child under 12 years of age for more than 5 days without consulting your doctor.
- If your child's cough comes back or is accompanied by a high temperature, rash or a headache that does not go away, stop using this medicine and talk to your doctor or pharmacist.

! If anyone has too much

If anyone has taken too much, contact a doctor or your nearest Accident and Emergency department (Casualty), taking this leaflet and pack with you.

! If you forget to give the medicine

You should only give this medicine as required following the dosage instructions above carefully. If you forget to give a dose, give the next dose when needed. **Do not** give a double dose.

4 Possible side-effects

Like all medicines, this medicine can cause side-effects, although not everybody gets them.

If your child experiences any of the following stop using the medicine and seek immediate help:

- Allergic reactions including skin rashes (which may be severe and include blistering and peeling), hives and itching.

Other effects that may occur include:

- Stomach pain or discomfort, diarrhoea, vomiting and nausea.

Reporting of side-effects

If your child experiences any side-effects, talk to your doctor, pharmacist or nurse. This includes any possible

side-effects not listed in this leaflet. You can also report side-effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side-effects, you can help provide more information on the safety of this medicine.

5 How to store this medicine

Do not store this product above 25°C.

Discard the bottle 4 months after opening, even if there is syrup remaining.

Store in the original package.

Keep the product out of the sight and reach of children.

Do not use after the end of the month shown as an expiry date on the packaging.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6 Contents of the pack and other information

What's in this medicine?

The active ingredient in 5 ml of Benylin Children's Chesty Coughs is: 50 mg Guaifenesin.

Other ingredients are: Glycerol, sorbitol liquid (E420), carmellose sodium, sodium citrate, sodium saccharin, sodium benzoate (E211), citric acid monohydrate, strawberry flavour (containing benzyl alcohol) and water.

What the medicine looks like

Benylin Children's Chesty Coughs is a clear colourless syrup, available in 125 ml, glass bottles.

Product Licence holder: McNeil Products Ltd, High Wycombe, Buckinghamshire, HP12 4EG, UK.

Manufacturer: Delpharm Orleans, 5 avenue de Concy, 45071 Orleans, Cedex 2, France.

This leaflet
Benylin is a

7 Information treating in children

It's normal for
Gradually the
Most colds
need to do
they get over
not bacterial
steps to help

- **If they are**
your child
with a par
doses for
under 2 m
3 months

- **For cough**
child cough
helps clear
nose. Give
drink.

- **To help v**
available t
blocked n

McNeil Pro

McN

© 2011